

## Key Facts

---

It is estimated that between 1 in 10,000 - 25,000 people in the population have Cornelia de Lange syndrome.

---

Most diagnoses are made based on the [clinical features](#) of the syndrome.

---

Cornelia de Lange syndrome is caused by a single [genetic change](#) affecting a single gene in each individual. However, the genetic change can be in different genes in different individuals.

---

Research studies have suggested that most individuals with Cornelia de Lange syndrome had a severe to profound [intellectual disability](#). Some individuals with Cornelia de Lange syndrome have mild intellectual disabilities.

---

Some of the [health difficulties](#) in Cornelia de Lange syndrome may be particularly important for understanding [behaviours that challenge](#).

---

Children and adults with Cornelia de Lange syndrome show higher levels of [compulsive-like behaviours](#) in comparison to other genetic syndromes.

---

While many children and adults meet the cut-off for [Autism Spectrum Disorder](#) (ASD) on standard assessments, the profile of ASD characteristics may be different in Cornelia de Lange syndrome.

---

Everyday living skills may be delayed among children with Cornelia de Lange syndrome but some do achieve a degree of independence in this area.

---

Many individuals with Cornelia de Lange syndrome experience [social anxiety](#) and are selectively mute.

---

Between 50-60% of individuals with Cornelia de Lange syndrome engage in some form of [self-injurious behaviour](#).

---

[Overactivity](#) seen in children and adults with Cornelia de Lange syndrome may be linked to pain and discomfort associated with gastro-intestinal difficulties.

---